

TM 1466 – RAPID-SENSITIVITY TEST BROTH

INTENDED USE

For antimicrobial susceptibility test.

PRODUCT SUMMARY AND EXPLANATION

The goal of an antimicrobial susceptibility test is to predict through an in vitro assessment the likelihood of successfully treating an infection with a particular antimicrobial agent. There are several continual or novel methods for performing antibacterial susceptibility testing. These include the disk diffusion test, broth microdilution, agar gradient and rapid automated instrument methods. Rapid Sensitivity Test Broth, which is used for antimicrobial susceptibility tests, is a semi-defined medium in which the mineral contents have been stabilized to give reproducible results. The thiamine and thymidine content is very low thus making it most suitable for testing antimicrobial activity of sulphonamides. However, some mutant strains which are totally dependent on thiamine and thymidine for their growth, will not grow in Rapid-Sensitivity Test Broth, due to very low levels of these compounds in the media as they are the naturally occurring antagonist of trimethoprim.

COMPOSITION

Ingredients	Gms / Ltr
Casein enzymic hydrolysate	11.000
Peptic digest of animal tissue	3.000
Dextrose	2.000
Sodium chloride	3.000
Starch, soluble	1.000
Disodium phosphate	2.000
Sodium acetate	1.000
Magnesium glycerophosphate	0.200
Calcium gluconate	0.100
Cobaltous sulphate	0.001
Cupric sulphate	0.001
Ferrous sulphate	0.001
Zinc sulphate	0.001
Manganous chloride	0.002
Menadione	0.001
Cyanocobalamin	0.001
L-Cysteine hydrochloride	0.020
L-Tryptophan	0.020
Pyridoxine hydrochloride	0.003



Calcium pantothenate	0.003
Nicotinamide	0.003
Biotin	0.0003
Thiamine hydrochloride	0.00004
Adenine	0.010
Guanine	0.010
Xanthine	0.010
Uracil	0.010

PRINCIPLE

This medium consists of Casein enzymic hydrolysate, peptic digest of animal tissue, dextrose, and vitamins provides nitrogen, carbon compounds and other essential growth nutrients.

INSTRUCTION FOR USE

- Dissolve 23.4 grams in 1000 ml purified/distilled water.
- Heat to boiling to dissolve the medium completely.
- Sterilize by autoclaving at 15 psi pressure (121°C) for 15 minutes.
- Mix well and dispense as desired.

QUALITY CONTROL SPECIFICATIONS

Appearance of Powder : Cream to yellow homogeneous free flowing powder.

Appearance of prepared medium : Basal medium: Light yellow ; After addition of 5%v/v laked blood : Red to chocolate coloured, Basal medium :clear to slightly opalescent ; After Addition: opalescent solution in tubes.

pH (at 25°C) : 7.4 ± 0.2

INTERPRETATION

Cultural characteristics observed after incubation.

Microorganism	ATCC	Inoculum (CFU/ml)	Growth	Incubation Temperature	Incubation Period
<i>Bacillus subtilis</i>	6633	50-100	Good-luxuriant	35-37°C	18-24 Hours
<i>Bacteroides vulgatus</i>	8482	50-100	Good-luxuriant	35-37°C	18-24 Hours
<i>Enterococcus faecalis</i>	29212	50-100	Good-luxuriant	35-37°C	18-24 Hours



<i>Salmonella</i> Typhimurium	14028	50-100	Good-luxuriant	35-37°C	18-24 Hours
<i>Staphylococcus</i> <i>aureus</i>	25923	50-100	Good-luxuriant	35-37°C	18-24 Hours
<i>Streptococcus</i> <i>pyogenes</i>	19615	50-100	Good-luxuriant	35-37°C	18-24 Hours

PACKAGING:

In pack size of 100 gm and 500 gm bottles.

STORAGE

Dehydrated powder, hygroscopic in nature, store in a dry place, in tightly-sealed containers between 25-30°C and protect from direct sunlight. Under optimal conditions, the medium has a shelf life of 4 years. When the container is opened for the first time, note the time and date on the label space provided on the container. After the desired amount of medium has been taken out replace the cap tightly to protect from hydration.













Product Deterioration: Do not use if they show evidence of microbial contamination, discoloration, drying or any other signs of deterioration.

DISPOSAL

After use, prepared plates, specimen/sample containers and other contaminated materials must be sterilized before discarding.

REFERENCES

1. Murray P. R., Baron J. H., Pfaller M. A., Jorgensen J. H. and Tenover F. C., (Ed.). 2003, Manual of Clinical Microbiology, 8th Ed., American Society for Microbiology, Washington, D.C.
2. Tanner E. I. and Bullin C. H., 1974, J. Clin. Path., 27:565.
3. Thomas M. and Bond L., 1973, Med. Lab. Technol., 30:277.
4. Barker J., Healing D., and Hutchinson J. G. P., 1972, J. Clin. Path., 25:1086
5. Ericsson H. M. and Sherris J. C., 1971, Acta. Pathol. Microbiol Scand Suppl., 217:1.
6. Garrod L. P. and Waterworth P. M., 1971, J. Clin. Path., 24:779.

 GMP Good Manufacturing Practices Certified	 IVD For In Vitro Diagnostic Use	 QTY. Quantity	 LOT/ B. NO. Lot / Batch Number	 REF Catalogue Number	 Manufacturer
 Temperature Unit	 EC REP Authorized Representative <small>MedNet GmbH Buckrose 10, 49163 Moenster, Germany</small>	 European Conformity	 QR Code	 Consults Instructions for Use	 Best Before

NOTE: Please consult the Material Safety Data Sheet for information regarding hazards and safe handling Practices.

***For Lab Use Only**
Revision: 08 Nov., 2019