

TM 2006 – BILE ESCULIN AZIDE AGAR, MODIFIED

INTENDED USE

For rapid, selective detection and enumeration of Enterococci and Group D Streptococci.

PRODUCT SUMMARY AND EXPLANATION

Group D Streptococci possess the group D lipoteichoic acid antigen in their cell walls. Group D species, are predominant normal inhabitants of the human gastrointestinal tract, are termed as faecal streptococci or Enterococci. The unique ability of Enterococci to split esculin was reported by Meyer and Schonfeld. Enterococci and group D streptococci hydrolyze esculin to esculetin and dextrose, which reacts with ferric citrate producing brownish black precipitate. The use of esculin hydrolysis in identification of Enterococci was first cited by Rochaix. However, other tests such as salt tolerance should be performed for identifying Enterococci. Modified Bile Esculin Azide Agar was formulated according to Isenberg et al, Swan, Facklam and Moody and Meyer and Schonfeld. They reported that esculin hydrolysis and bile tolerance permit the isolation and identification of group D streptococci in 24 hours.

COMPOSITION

Ingredients	Gms / Ltr
Tryptone	17.000
Yeast extract	5.000
Peptone	3.000
Oxgall	10.000
Esculin	1.000
Ferric ammonium citrate	0.500
Sodium chloride	5.000
Sodium citrate	1.000
Sodium azide	0.250
Agar	15.000

PRINCIPLE

Tryptone, proteose peptone and beef extract serves as sources of carbon, nitrogen, amino acids, vitamins and essential growth nutrients. oxgall and sodium azide inhibits most of the other accompanying bacteria. Esculin in the medium is hydrolyzed to esculetin and dextrose. Esculetin reacts with ferric citrate to form a dark brown or black complex, visualized as a zone of black precipitate around the colonies.

INSTRUCTION FOR USE

- Dissolve 56.65 grams in 1000 ml distilled water.
- Heat to boiling to dissolve the medium completely.
- Sterilize by autoclaving at 15 psi pressure (121°C) for 15 minutes. Cool to 45-50°C.
- Mix well and pour into sterile Petri plates.

QUALITY CONTROL SPECIFICATIONS



Appearance of Powder : Cream to brownish yellow homogeneous free flowing powder.
Appearance of prepared medium : Medium amber coloured clear to slightly opalescent solution with a bluish tinge forms in Petri plates.
pH (at 25°C) : 7.1±0.2

INTERPRETATION

Cultural characteristics observed after incubation.

Microorganism	ATCC	Inoculum (CFU/ml)	Growth	Recovery	Esculin hydrolysis	Incubation Temperature	Incubation Period
<i>Enterococcus faecalis</i>	29212	50-100	Luxuriant	≥70%	Positive reaction, blackening of medium around the colony	35-37°C	18-24 Hours
<i>Escherichia coli</i>	25922	50-100	Inhibited	0%	-	35-37°C	18-24 Hours
<i>Staphylococcus aureus</i>	25923	50-100	Good	40-50%	Negative reaction	35-37°C	18-24 Hours
<i>Proteus mirabilis</i>	25933	50-100	Good	40-50%	Negative reaction	35-37°C	18-24 Hours
<i>Streptococcus pyogenes</i>	19615	50-100	None-poor	0-10%	Negative reaction	35-37°C	18-24 Hours

PACKAGING:

In pack size of 500 gm bottles.

STORAGE

Dehydrated powder, hygroscopic in nature, store in a dry place, in tightly-sealed containers between 25-30°C and protect from direct sunlight. Under optimal conditions, the medium has a shelf life of 4 years. When the container is opened for the first time, note the time and date on the label space provided on the container. After the desired amount of medium has been taken out replace the cap tightly to protect from hydration.

Product Deterioration: Do not use if they show evidence of microbial contamination, discoloration, drying or any other signs of deterioration.

DISPOSAL













After use, prepared plates, specimen/sample containers and other contaminated materials must be sterilized before discarding.

REFERENCES

- Koneman E. W., Allen S. D., Janda W. M., Schreckenberger P. C., Winn W. C. Jr., 1992, Colour Atlas and Textbook of Diagnostic Microbiology, 4 th Ed., J. B. Lippincott Company
- Meyer and Schonfeld, 1926, Zentralbl. Bakteriologie, Parasitenkunde, Infektionskrankheiten und Hygiene, 99:402.
- MacFaddin J. F., 1985, Media for Isolation-Cultivation-Identification-Maintenance of Medical Bacteria, Vol. I, Williams and Wilkins, Baltimore.
- Rochaix, 1924, Compt. Rend. Soc. Biol., 90:771.
- Facklam R., 1973, Appl. Microbiol., 26:138.
- Swan, 1954, J. Clin. Pathol., 7:160.



7. Facklam R., 1972, Appl. Microbiol., 23:1131.
8. Facklam R. R and Moody M. D., 1970, Appl. Microbiol., 20(2):245.
9. Edberg S. C., Pittman S., and Singer J. M., 1977, J. Clin. Microbiol., 6:111.
10. Isenberg, 1970, Clin. Lab. Forum, July.
11. Murray P. R., Baron E. J., Jorgensen J. H., Pfaller M. A., Tenover F. C., Tenover F. C. (Eds.), 8th Ed., 2003, Manual of Clinical Microbiology, ASM, Washington, D.C.

 GMP Good Manufacturing Practices Certified	 IVD For In Vitro Diagnostic Use	 QTY. Quantity	 LOT/ B. NO. Lot / Batch Number	 REF Catalogue Number	 Manufacturer
 Temperature Unit	 EC REP MedNet GmbH Birkstrasse 10, 48163 Muenster, Germany Authorized Representative	 European Conformity	 QR Code	 Consults Instructions for Use	 Best Before

NOTE: Please consult the Material Safety Data Sheet for information regarding hazards and safe handling Practices.

***For Lab Use Only**
Revision: 08 Nov., 2019