1

f (0) in 🔰



TM 648 – ALGAE CULTURE BROTH

INTENDED USE

For isolation and cultivation of algae from soil and water.

PRODUCT SUMMARY AND EXPLANATION

Algae (singular alga) encompass several groups of relatively simple living aquatic organisms that capture light energy through photosynthesis, using it to convert inorganic substances into organic matter. Algae range from single-cell organisms to multicellular organisms, some with fairly complex differentiated form and (if marine) called seaweeds. Algae are usually found in damp places or water bodies and thus are common in terrestrial as well as aquatic environments. Various algae play significant roles in aquatic ecology. Algae are used by humans in a number of ways. Because many species are aquatic and microscopic, they are cultured in clear tanks or ponds and either harvested or used to treat effluents pumped through ponds. Algae Culture Broth is recommended for the isolation and cultivation of algae from soil, water and sewage. Algae Culture Broth is used to prepare the inoculum for the bioassay of algicidal chemicals. Algae Culture Broth is similar in composition to Algae Culture Agar, except the agar.

COMPOSITION

Ingredients	Gms / Ltr	
Sodium nitrate	1.000	
Dipotassium hydrogen phosphate	0.250	
Magnesium sulphate	0.513	
Ammonium chloride	0.050	
Calcium chloride	0.058	
Ferric chloride	0.003	

PRINCIPLE

The medium provides all necessary nutrients for good growth of Algae but does not provide for other than minimal growth of bacteria and fungi. The salt provides nutrients as well as balances the osmotic pressure. Agar acts as a solidifying agent.

INSTRUCTION FOR USE

- Dissolve 1.87 grams in 1000 ml purified / distilled water.
- Heat if necessary to dissolve the medium completely.
- Dispense in tubes or flasks as desired. Sterilize by autoclaving at 15 psi pressure (121°C) for 15 minutes.

QUALITY CONTROL SPECIFICATIONS

Appearance of Powder	: White to light yellow homogeneous free flowing powder.
Appearance of prepared medium	: White coloured clear to slightly opalescent solution in tubes.
pH (at 25°C)	: 7.0±0.2

INTERPRETATION

Cultural characteristics observed after incubation.

A- 902A, RIICO Industrial Area, Phase III, Bhiwadi-301019.



Microorganism	ATCC	lnoculum (CFU/ml)	Growth	Incubation Temperature	Incubation Period
Chlorella pyrenoidosa	50476	50-100	Good-luxuriant	20-25°C	1 Week

PACKAGING:

In pack size of 100 gm and 500 gm bottles.

STORAGE

Dehydrated powder, hygroscopic in nature, store in a dry place, in tightly-sealed containers between 25-30°C and protect from direct sunlight. Under optimal conditions, the medium has a shelf life of 4 years. When the container is opened for the first time, note the time and date on the label space provided on the container. After the desired amount of medium has been taken out replace the cap tightly to protect from hydration.

Product Deterioration: Do not use if they show evidence of microbial contamination, discoloration, drying or any other signs of deterioration.

DISPOSAL

After use, prepared plates, specimen/sample containers and other contaminated materials must be sterilized before discarding.


REFERENCES

1. Baird R.B., Eaton A.D., and Rice E.W., (Eds.), 2015, Standard Methods for the Examination of Water and Wastewater, 23rd ed., APHA, Washington, D.C.

2.Guiry M. D. and Blunden G., (Ed.), 1991, Seaweed Resources in Europe: Uses and Potential. John Wiley and Sons Ltd.

3. Lembi C. A. and Waaland J. R., (Ed.), Algae and Human Affairs, 1988, Cambridge University Press.

4. Subba Rao N. S., 1977, Soil Microorganisms and Plant Growth, Oxford and IBH Publishing Co., New Delhi.



NOTE: Please consult the Material Safety Data Sheet for information regarding hazards and safe handling Practices. *For Lab Use Only Revision: 08 Nov., 2019

