

TMT 035 – VIRAL TRANSPORT MEDIUM W/ GLASS BEADS

INTENDED USE

For collection and transport of clinical specimen for recovery of viral agents.

PRODUCT SUMMARY AND EXPLANATION

A Viral Transport Medium is intended for the collection and transport of clinical specimens containing viruses, chlamydiae, mycoplasmas or ureaplasmas from the collection site to the testing laboratory. Viral Transport Medium is designed to maintain viral viability and transport viruses in active form for isolation. The peculiar design of the flocked swab ensures optimal elution of the specimen into the transport medium. The viral transport medium contains essential buffers and antibiotics required for maintaining the viability of the viruses during transport. The medium is also recommended by CDC and WHO for collection and transport of Coronavirus. Glass beads in each tube aid in the release and dispersion of patient material and viral particles from the swab.

COMPOSITION

Components	Composition
Viral Transport Medium*	Proprietary
Glass Beads	-

^{*3}ml medium in 10-15 ml tube

PRINCIPLE

The viral transport medium consists of Hanks Balanced Salt Solution modified and enriched with bovine serum albumin, cysteine, gelatin, sucrose and glutamic acid. The pH is quenched with buffer HEPES. Phenol red is used as a pH indicator. Vancomycin, amphotericin B and colistin have been added to the medium to inhibit the proliferation of competing bacteria and yeasts. The medium is isotonic and lacks toxicity to the mammalian host cells. The presence of sucrose acts as a cryoprotectant that facilitates the viruses and chlamydia if samples are frozen (-70 ° C) for long storage.

QUALITY CONTROL SPECIFICATIONS

Appearance Orange-red colour, clear solution

pH (at 25°C) 7.3 ± 0.2

Sterility Check Passes release criteria

INSTRUCTION FOR USE

Label the sample correctly with name of patient, time and date of collection. Inoculate the sample and transport the samples immediately to the laboratory for processing.

TRANSPORTATION OF SAMPLE:

To maintain optimum viability, transport the specimen to the laboratory as soon as possible. Best recovery is obtained when specimens are refrigerated at 2-8°C or kept on wet ice following collection and while in transit. If there will be long delay before processing, it is suggested that specimen should be frozen at -70°C.

STORAGE AND SHELF LIFE:

The viral transport medium should be stored at 15-30°C before sample collection and 2-8°C after sample collection. Use before the expiry date.

PRECAUTIONS













PRODUCT DATA SHEET

- Isolation of viruses will largely depend on proper specimen collection, timing of sample collection and processing
- Do not use the product if, (i) there is change in the color of the medium, (ii) there is evidence of leakage and (iii) 2. there are other signs of deterioration.
- 3. Specimen collection should be done in the acute phase of illness.
- 4. Avoid repeated freeze-thaw of collected samples.
- To maintain infectivity of viruses, it is important that temperature be properly maintained for sample collection 5. to processing.
- 6. It is recommended to refer to standard procedures and published protocols for sample collection and processing.























IVD For In Vitro Diagnostic Use

NOTE: Please consult the Material Safety Data Sheet for information regarding hazards and safe handling Practices.

*For Lab Use Only

Revision: 29th March., 2022







