

TMV 322 - MacCONKEY BROTH W/ NEUTRAL RED (VEG.)

INTENDED USE

For selective enrichment and enumeration of coliforms from water.

PRODUCT SUMMARY AND EXPLANATION

This medium is a modification of MacConkey Broth with Neutral Red which is considered as a standard medium for the primary isolation as well as presumptive identification of coliform-aerogenes group of organisms in food and water. The medium contains neutral red as an indicator.

COMPOSITION

Ingredients	Gms / Ltr
Veg peptone	23.000
Lactose	10.000
Synthetic detergent	2.000
Sodium chloride	5.000
Neutral red	0.075

PRINCIPLE

Veg Peptone provides necessary nitrogen source. Lactose serves as the fermentable carbohydrate source. Sodium chloride maintains the osmotic balance of the cells. The selective action of these media is attributed to the presence of bile salts, which are inhibitory to most species of gram-positive bacteria. Gram-negative bacteria usually grow well on these media and are differentiated by their ability to ferment lactose. The colour change of the medium shown by lactose-fermenters is due to production of acid from lactose and a subsequent colour change of the indicator dye when the pH of the media falls below 6.8. Lactose non-fermenting strains, such as *Shigella* and *Salmonella* do not alter the appearance of the medium. The medium turns pink in case of lactose fermenters and yellow in case of non-lactose-fermenters, due to neutral red. MacConkey Broth, which contains neutral red as an indicator is considered as a standard medium for the primary isolation as well as presumptive identification of coliform-aerogenes group of organisms in food and water.

INSTRUCTION FOR USE

- Dissolve 40.07 grams in 1000 ml purified/distilled water.
- Heat if necessary to dissolve the medium completely.
- Distribute into tubes with inverted Durham's tubes.
- Sterilize by autoclaving at 15 psi pressure (121°C) for 15 minutes.
- Cool the tubes before inoculation.

QUALITY CONTROL SPECIFICATIONS

Appearance of Powder	: Yellow coloured may have slightly greenish tinge, homogeneous, free flowing powder.
Appearance of prepared medium	: Red coloured, clear solution without any precipitate.
pH (at 25°C)	: 7.4±0.2

INTERPRETATION

Cultural characteristics observed after an incubation.



Microorganism	ATCC	Inoculum (CFU/ml)	Growth	Acid	Gas	Incubation Temperature	Incubation Period
<i>Enterobacter aerogenes</i>	13048	50-100	Luxuriant	Positive reaction	Positive reaction	35-37°C	18-24 Hours
<i>Escherichia coli</i>	25922	50-100	Luxuriant	Positive reaction	Positive reaction	35-37°C	18-24 Hours
<i>S. serotype Cholerasuis</i>	12011	50-100	Fair to good	-	-	35-37°C	18-24 Hours
<i>Staphylococcus aureus</i>	25923	$\geq 10^3$	Inhibited	-	-	35-37°C	18-24 Hours

PACKAGING:

In pack size of 100 gm and 500 gm bottles.

STORAGE

Dehydrated powder, hygroscopic in nature, store in a dry place, in tightly-sealed containers between 25-30°C and protect from direct sunlight. Under optimal conditions, the medium has a shelf life of 4 years. When the container is opened for the first time, note the time and date on the label space provided on the container. After the desired amount of medium has been taken out replace the cap tightly to protect from hydration.

Product Deterioration: Do not use if they show evidence of microbial contamination, discoloration, drying or any other signs of deterioration.













DISPOSAL

After use, prepared plates, specimen/sample containers and other contaminated materials must be sterilized before discarding.

REFERENCES

- Greenberg A. E., Trussell R. R. and Clesceri L. S. (Eds.), Standard Methods for the Examination of Water and Wastewater, 1985, 16th ed., A.P.H.A., Washington, D.C.
- Rappaport F. and Henigh E., 1952, J. Clin. Path., 5:361.
- International Organization for Standardization (ISO), 1990, Draft ISO/DIS 9308-2.
- Harrigan W.F. and McCance M.E. (Eds.), 1976, Laboratory Methods in Food and Dairy Microbiology, Academic Press, London.
- Holt, Harris and Teague, 1916, J. Infect. Dis., 18:596.
- MacConkey, 1900, The Lancet, ii:20.
- MacConkey, 1905, J. Hyg., 5:333.
- Speck M. (Ed.), 1985, Compendium of Methods for the Microbiological Examination of Foods, 2nd ed., APHA, Washington, D.C.
- Greenberg A. E., Clesceri L. S. and Eaton A. D., (Eds.), 1992, Standard Methods for the Examination of Water and Wastewater, 18th ed., APHA, Washington, D.C.
- Marshall R. (Ed.), 1992, Standard Methods for the Examination of Dairy Products, 16th ed., APHA, Washington, D.C.
- Karmali M.A., Petric M., Lim C., et al, 1985, J. Infect. Dis., 151:775.
- Lior H. and Borczyk A., 1987, Lancet, i:333.
- MacFaddin J., 1985, Media for Isolation-Cultivation-Identification- Maintenance of Medical Bacteria, Vol. I, Williams and Wilkins, Baltimore.



 GMP Good Manufacturing Practices Certified	 IVD For In Vitro Diagnostic Use	 QTY. Quantity	 LOT/ B. NO. Lot / Batch Number	 REF Catalogue Number	 Manufacturer
 Temperature Unit	 EC REP Authorized Representative <small>MedNet GmbH Buckenhof 10 48143 Aachen, Germany</small>	 European Conformity	 QR Code	 Consults instructions for Use	 Best Before

NOTE: Please consult the Material Safety Data Sheet for information regarding hazards and safe handling Practices.

***For Lab Use Only**
Revision: 08 Nov., 2019