

## TS 005 – POTASSIUM TELLURITE 1% (1 ml /vl)

### INTENDED USE

For the selective isolation of Staphylococci and *Corynebacteria*.

### COMPOSITION

Ingredients	Concentration
Potassium tellurite Concentrate	1.100ml

(To achieve 1% solution dilute the contents in 8.9 ml sterile distilled water)

### Applied in Medium

TM 636 - AGAR MEDIUM O (BAIRD PARKER AGAR BASE) (as per USP/EP/BP)  
 TM 670 - BAIRD STAPHYLOCOCCUS ENRICHMENT BROTH BASE  
 TM 411 - CHOLERA MEDIUM BASE  
 TM 707 - CYSTINE TELLURITE AGAR BASE  
 TM 1165 - DEXTROSE PROTEOSE PEPTONE AGAR BASE  
 TM 984 - DIPHTHERIA VIRULENCE AGAR BASE  
 TM 221 - MITIS SALIVARIUS AGAR BASE  
 TMV 221 - MITIS SALIVARIUS AGAR BASE (VEG.)  
 TM 1048 - MONSUR MEDIUM BASE  
 TM 235 - MYCOPLASMA BROTH BASE W/ CV (PPLO BROTH BASE W/ CV)  
 TM 438 - TPEY AGAR BASE  
 TM 879 - TELLURITE BLOOD AGAR BASE  
 TM 439 - TELLURITE GLYCINE AGAR BASE  
 TM 1109 - TRYPTONE TELLURITE AGAR BASE  
 TM 395 - VOGEL-JOHNSON AGAR BASE W/O TELLURITE (V. J. AGAR BASE)  
 TMV 395 - VOGEL JOHNSON AGAR BASE W/O TELLURITE (V. J. AGAR BASE) (VEG.)  
 TM 1722 - VOGEL JOHNSON AGAR MEDIUM (as per USP)  
 TM 1820 - VOGEL JOHNSON AGAR MEDIUM (as per IP)  
 TM 2121 - CHROMOGENIC EC O157:H7 AGAR, MODIFIED  
 TM 2420 - VOGEL JOHNSON AGAR BASE W/ 1.5% AGAR  
 TMH 119 - BAIRD PARKER AGAR BASE (as per EP/IP/BP)







### INSTRUCTION FOR USE

Warm up the refrigerated contents of one vial to room temperature. Add aseptically 8.9 ml sterile distilled water, mix well and add in sterile, molten, cooled (45-50°C) TM 636 - AGAR MEDIUM O (BAIRD PARKER AGAR BASE) (as per USP/EP/BP) / TMH 119 - BAIRD PARKER AGAR BASE (as per EP/IP/BP) / TM 670 - BAIRD STAPHYLOCOCCUS ENRICHMENT BROTH BASE / TM 395 - VOGEL-JOHNSON AGAR BASE W/O TELLURITE (V. J. AGAR BASE) / TMV 395 - VOGEL JOHNSON AGAR BASE W/O TELLURITE (V. J. AGAR BASE) (VEG.) / TM 1722 - VOGEL JOHNSON AGAR MEDIUM (as per USP) / TM 1820 - VOGEL JOHNSON AGAR MEDIUM (as per IP) / TM 235 - MYCOPLASMA BROTH BASE W/ CV (PPLO BROTH BASE W/ CV) / TM 438 - TPEY AGAR BASE / TM 439 - TELLURITE GLYCINE AGAR BASE / TM 411 - CHOLERA MEDIUM BASE / TM 707

- CYSTINE TELLURITE AGAR BASE / TM 1165 - DEXTROSE PROTEOSE PEPTONE AGAR BASE / TM 984 - DIPHTHERIA VIRULENCE AGAR BASE / TM 1109 - TRYPTONE TELLURITE AGAR BASE / TM 879 - TELLURITE BLOOD AGAR BASE / TM 221 - MITIS SALIVARIUS AGAR BASE / TMV 221 - MITIS SALIVARIUS AGAR BASE (VEG.) / TM 1048 - MONSUR MEDIUM BASE / TM 2121 - CHROMOGENIC EC O157:H7 AGAR, MODIFIED / TM 2420 - VOGEL JOHNSON AGAR BASE W/ 1.5% AGAR. Mix well and dispense in sterile Petri plates or tubes.

**STORAGE**

Vials should be stored in sealed container at 2-8°C.

<b>GMP</b> Good Manufacturing Practices Certified	<b>IVD</b> For In Vitro Diagnostic Use	<b>QTY.</b> Quantity	<b>LOT/ B. NO.</b> Lot / Batch Number	<b>REF</b> Catalogue Number	 Manufacturer
 Temperature Unit	<b>EC REP</b> Authorized Representative <small>MedWor GmbH Sachsenstr. 10 48147 Münster, Germany</small>	 European Conformity	 QR Code	 Consults Instructions for Use	 Best Before

**NOTE:** Please consult the Material Safety Data Sheet for information regarding hazards and safe handling Practices.

**\*For Lab Use Only**  
Revision: 4 March., 2022