

## TS 011 – STREPTO SUPPLEMENT

### INTENDED USE

For selective cultivation *Streptococcs* species.

### COMPOSITION

Ingredients	Concentration
Nalidixic acid	3.750mg
Neomycin sulphate	1.060mg
Polymyxin B sulphate	8500Units

(per vial sufficient for 500 ml medium)

### Applied in Medium


- TM 041 - BLOOD AGAR BASE No. 2
- TM 1842 - BLOOD AGAR BASE NO. 2 (ISO 11290-1:2017)
- TM 2011 - BLOOD AGAR BASE NO. 2 WITH 1.2% AGAR
- TM 071 - COLUMBIA BLOOD AGAR BASE
- TMV 071 - COLUMBIA BLOOD AGAR BASE (VEG.)
- TM 2044 - COLUMBIA BLOOD AGAR BASE W/ 1% AGAR

### INSTRUCTION FOR USE

Rehydrate the contents of 1 vial aseptically with 2-5 ml of sterile distilled water. Aseptically add the contents to 500 ml of sterile, molten, cooled (45-50°C) TM 071 - COLUMBIA BLOOD AGAR BASE / TMV 071 - COLUMBIA BLOOD AGAR BASE (VEG.) / TM 2044 - COLUMBIA BLOOD AGAR BASE W/ 1% AGAR / TM 041 - BLOOD AGAR BASE No. 2 / TM 1842 - BLOOD AGAR BASE NO. 2 (ISO 11290-1:2017) / TM 2011 - BLOOD AGAR BASE NO. 2 WITH 1.2% AGAR. The medium may be enriched with 5-7% v/v sterile defibrinated blood. Mix gently and pour into sterile Petri plates.

### STORAGE

Vials should be stored in sealed container at 2-8°C.



**NOTE:** Please consult the Material Safety Data Sheet for information regarding hazards and safe handling Practices.

**\*For Lab Use Only**  
Revision: 3 March., 2022

