

TS 035 – FRASER SELECTIVE SUPPLEMENT

INTENDED USE

For selective isolation and enrichment of *Listeria monocytogenes* from food and environmental samples.

COMPOSITION

Ingredients	Concentration
Acriflavine hydrochloride	25mg
Nalidixic acid	20mg

(per vial sufficient for 1000 ml medium)

Applied in Medium

TM 113 - FRASER SECONDARY ENRICHMENT BROTH BASE

TM 1472 - FRASER BROTH BASE (ISO 11290-1: 2017)

TM 1937 - HALF FRASER BROTH BASE (ISO 11290-1:2017)


TM 2092 - FRASER BROTH BASE

INSTRUCTION FOR USE

Rehydrate the contents of 1 vial aseptically with 10 ml of sterile distilled water. Mix well and aseptically add to 990 ml of sterile, cooled (45-50°C) TM 113 - FRASER SECONDARY ENRICHMENT BROTH BASE / TM 1472 - FRASER BROTH BASE (ISO 11290-1: 2017) / TM 1937 - HALF FRASER BROTH BASE (ISO 11290-1:2017) / TM 2092 - FRASER BROTH BASE. Mix well and dispense as desired.

STORAGE

Vials should be stored in sealed container at 2-8°C.



NOTE: Please consult the Material Safety Data Sheet for information regarding hazards and safe handling Practices.

***For Lab Use Only**
Revision: 28 Feb., 2022