

## TS 051 – NOVOBIOCIN SUPPLEMENT

### INTENDED USE

For presumptive identification of *Aeromonas hydrophila*.

### COMPOSITION

Ingredients	Concentration
Novobiocin	5mg

(per vial sufficient for 1000 ml medium)

### Applied in Medium

TM 1400 - MULLER KAUFFMAN TETRATHIONATE BROTH BASE

TM 1127 - SEMISOLID RAPPAPORT VASSILIADIS MEDIUM, MODIFIED

TM 546 - SOYABEAN BILE BROTH BASE

TM 2056 - DOUBLE MODIFIED LYSINE IRON AGAR BASE













TM 2301 - RS MEDIUM BASE

### INSTRUCTION FOR USE

Rehydrate the contents of 1 vial aseptically with 2 ml of sterile distilled water and aseptically add to 1000 ml of sterile, molten, cooled (45-50°C) TM 1400 - MULLER KAUFFMAN TETRATHIONATE BROTH BASE / TM 1127 - SEMISOLID RAPPAPORT VASSILIADIS MEDIUM, MODIFIED / TM 546 - SOYABEAN BILE BROTH BASE / TM 2056 - DOUBLE MODIFIED LYSINE IRON AGAR BASE / TM 2301 - RS MEDIUM BASE. Mix well and pour into sterile petri plates.

### STORAGE

Vials should be stored in sealed container at 2-8°C.

 GMP Good Manufacturing Practices Certified	 IVD For In Vitro Diagnostic Use	 QTY. Quantity	 LOT/ B. NO. Lot / Batch Number	 REF Catalogue Number	 Manufacturer
 Temperature Unit	 EC REP Authorized Representative <small>Muller GmbH Bachstrasse 10 41143 Aachen, Germany</small>	 European Conformity	 QR Code	 Consults Instructions for Use	 Best Before

**NOTE:** Please consult the Material Safety Data Sheet for information regarding hazards and safe handling Practices.

\*For Lab Use Only  
Revision: 4 March., 2022