

TS 054 – S.F. P. SUPPLEMENT (Perfringens S.F.P. Supplement)

INTENDED USE

for selective isolation of *Clostridium perfringens*.

COMPOSITION

Ingredients	Concentration
Kanamycin sulphate	6mg
Polymyxin B	15000IU

(per vial sufficient for 500 ml medium)

Applied in Medium

TM 615 - PERFRINGENS AGAR BASE (T.S.C./S.F.P. AGAR BASE)

TMV 615 - PERFRINGENS AGAR BASE (T.S.C./S.F.P. AGAR BASE) (VEG.)

TM 846 - S.F.P. AGAR BASE


TM 615 - TRYPTOSE SULPHITE CYCLOSERINE AGAR BASE

INSTRUCTION FOR USE

Rehydrate the contents of 1 vial aseptically with 2 ml sterile distilled water. Mix well and aseptically add it to 475 ml of sterile, molten, cooled (45-50°C) TM 615 - PERFRINGENS AGAR BASE (T.S.C./S.F.P. AGAR BASE) / TMV 615 - PERFRINGENS AGAR BASE (T.S.C./S.F.P. AGAR BASE) (VEG.) / TM 615 - TRYPTOSE SULPHITE CYCLOSERINE AGAR BASE / TM 846 - S.F.P. AGAR BASE along with 25 ml of sterile Egg Yolk Emulsion. Mix well and pour into sterile petri plates.

STORAGE

Vials should be stored in sealed container at 2-8°C.



NOTE: Please consult the Material Safety Data Sheet for information regarding hazards and safe handling Practices.

***For Lab Use Only**
Revision: 3 March., 2022