

TS 148 – LISTERIA CONFIRMATORY SUPPLEMENT B

INTENDED USE

For isolation and enumeration of *Listeria monocytogenes*.

COMPOSITION

Ingredients	Concentration
Nalidixic acid	10mg
Cephatazidime	10mg
Amphoterecin	5mg

(Per vial sufficient for 500 ml medium)

Applied in Medium













TM 1375 - LISTERIA CONFIRMATORY AGAR BASE

INSTRUCTION FOR USE

Aseptically rehydrate 1 vial contents to 10ml of sterile distilled water and add to 500ml of sterile molten medium (cooled 45-50°C) TM 1375 - LISTERIA CONFIRMATORY AGAR BASE. Mix well and pour into the Petri plates. Store the prepared media plates inverted at 4-8°C for not more than 7 days.

STORAGE

Vials should be stored in sealed container at 2-8°C.

 GMP Good Manufacturing Practices Certified	 IVD For In Vitro Diagnostic Use	 QTY. Quantity	 LOT/ B. NO. Lot / Batch Number	 REF Catalogue Number	 Manufacturer
 Temperature Unit	 EC REP Authorized Representative <small>MedWor GmbH Buckhorn 16 40112 Mönchen, Germany</small>	 European Conformity	 QR Code	 Consults Instructions for Use	 Best Before

NOTE: Please consult the Material Safety Data Sheet for information regarding hazards and safe handling Practices.

***For Lab Use Only**
Revision: 11 March., 2022