

TS 296 – MUG SUPPLEMENT (50 MG PER VIAL)

INTENDED USE

For measuring b - glucuronidase activity, for rapid and sensitive identification of *Escherichia coli*.

COMPOSITION

Ingredients	Concentration
4-Methylumbeliferyl b-D-Glucuronide(MUG)	50mg

(Per vial sufficient for 500/1000 ml medium)

Applied in Medium













TM 2186 - M-FC BASAL MEDIUM

INSTRUCTION FOR USE

Rehydrate the contents of one vial aseptically with 5 ml methanol. One vial is sufficient for 500 ml agar media or 1000 ml of broth media. Add the contents of 1 vial to 1000 ml of sterile, molten, cooled (45-50°C) TM 2186 - M-FC BASAL MEDIUM. Mix well and pour into sterile petri plates.

STORAGE

Vials should be stored in sealed container at 2-8°C.

 GMP Good Manufacturing Practices Certified	 IVD For In Vitro Diagnostic Use	 QTY. Quantity	 LOT/ B. NO. Lot / Batch Number	 REF Catalogue Number	 Manufacturer
 Temperature Unit	 EC REP Authorized Representative	 European Conformity	 QR Code	 Consults Instructions for Use	 Best Before

NOTE: Please consult the Material Safety Data Sheet for information regarding hazards and safe handling Practices.

*For Lab Use Only
Revision: 8 March., 2022